



episil[®]
ORAL LIQUID

For fast and effective oral pain relief

*“It gave me the relief I needed to be able
to start eating properly.”*

Patient, UK

BTChealth.



episil[®] oral liquid

- ✓ Rapid pain relief, within 5 minutes^{1,2}
- ✓ Long-lasting effect, for up to 8 hours¹
- ✓ Ready-to-use, pocket-sized device³
- ✓ Developed in Sweden based on the patented and award-winning FluidCrystal[®] technology⁴

What is oral mucositis?

Oral mucositis is an inflammatory condition in the mouth. It is a frequent side effect of chemo- and radiotherapy. The condition causes redness, mouth sores and ulcers that can be very painful, making it difficult to eat, drink and speak.

What is episil[®] oral liquid?

episil[®] oral liquid is used for the relief of oral pain. When applied in the mouth, episil[®] oral liquid forms a bioadhesive film that protects and soothes oral lesions, thereby providing rapid and significant oral pain relief that lasts for up to 8 hours.

Content

Glycerol dioleate, Phosphatidylcholine (soy lecithin), Ethanol, Propylene glycol, Polysorbate 80 and Peppermint oil

Dosage

1-3 pump strokes, 2-3 times per day or as needed

Storage

✓ Keep out of reach and sight of small children

✓ Store in the original packaging at room temperature below 25°C

How and when to use episil® oral liquid?

Use episil® oral liquid for rapid and effective pain relief.



Remove the protective cap from the metering pump.



Prior to the first use, prime the pump mechanism by firmly pumping into a tissue until an even stream of liquid is delivered.



Firmly press the pump with your thumb and apply the stream of liquid to affected areas in the mouth 1-3 times.*



Distribute in the mouth by using the tongue. The protective film will form within a few minutes.



Repeat the treatment 2-3 times a day, or as needed. episil® oral liquid can also be applied prior to eating.

*If there is an abundant amount of mucus present, then it is recommended to rinse the mouth with water before using episil® oral liquid.

Possible side effects

There are no known risks or side effects. However, episil® oral liquid contains small amounts of ethanol (alcohol), which may cause a short, temporary burning sensation when applied to affected areas in the mouth. You should not use the product if you are allergic to any of its components. If you notice any side effects of episil® oral liquid, discontinue the treatment and contact your doctor. For more information about episil® oral liquid, please consult the package insert.

How to obtain episil® oral liquid

Ask your doctor or nurse about treatment with episil® oral liquid.

episil® oral liquid is available from your local pharmacy.
If it is not yet in stock, ask the pharmacist to order it for you.

The order code for episil® 10 mL is Symbion PDE 628239

www.btchealth.com.au

References

1. Hadjieva, T., Cavallin-Ståhl, E., Linden, M. and Tiberg, F. (2014) *Treatment of oral mucositis pain following radiation therapy for head-and-neck cancer using a bioadhesive barrier-forming lipid solution*. Support Care Cancer. 22(6):1557–1562. **2.** Seidenspinner, I., Adamietz, I. A., Strohm, G. L. and Tiberg, T. (2015) *Effects of episil® oral liquid in cancer patients with oral mucositis: an observational study*. Poster presented at the Multinational Association of Supportive Care in Cancer, 25–27 June 2015, Copenhagen, Denmark. **3.** Camurus. (2015) episil® package leaflet. **4.** FluidCrystal® was awarded the "Best innovation in formulation" prize at the CPhI Worldwide in 2013. <https://www.prnewswire.com/news-releases/cphi-pharma-awards-showcase-industry-drivers-229079101.html>
*Data on file.

BTC health.

episil® oral liquid is a registered trademark and is developed by Camurus AB, Ideon Science Park, SE-223 70, Lund, Sweden.
episil® oral liquid is distributed by BTC Speciality Health Pty Ltd (a wholly owned company of BTC Health) under licence from Camurus AB.
BTC Speciality Health, Suite 201, 697 Burke Road, Camberwell VIC 3124, Australia.
Website: btchealth.com.au. Queries: queries@btchealth.com.au.
Phone: Australia 1800 100 282 | New Zealand: 0800 567 370.
8234BTC.